



Identifying Providers Qualified to Manage the Longitudinal Treatment of Patients with HIV Infection and Resources to Support Quality HIV Care

Revised: March 2013

This document was developed to help health systems and third-party payers identify physicians who are best qualified to manage the care of HIV-infected patients and to compile clinical tools, standards and resources to promote high quality HIV care. The suggested standards are intended for medical providers managing the care of HIV-infected patients in an out-patient or clinic setting.

The criteria recommended below are based on a large body of evidence that indicates that, regardless of a physician’s specialty training, the two best predictors of high quality, cost effective HIV care are patient management experience and ongoing professional development through HIV-related continuing medical education.

Qualified HIV Physician Criteria

Category	Criteria
Patient Management	Management of at least 25 HIV-infected patients in the preceding 36 months.
Continuing Medical Education	At least 40 hours of HIV-related continuing medical education ¹ in the preceding 36 months, earning a minimum of 10 hours per year.
Board Certification Or Significant Clinical Experience	Board certification or equivalent in one or more medical specialties or subspecialties recognized by the American Board of Medical Specialties or the American Osteopathic Association is preferred. Significant clinical and professional experience in HIV medicine, defined as a minimum of at least five years, should be considered in the absence of board certification.

Infectious Diseases Physicians

Many HIV physicians are trained in infectious diseases (ID), but not all ID physicians continue to provide HIV care after training. Recently trained or recertified ID physicians should be considered qualified HIV physicians within 36 months of completing certification or recertification. However, ID physicians continuing to provide longitudinal care for HIV-infected patients should meet the criteria above

¹Completion of HIV-related Maintenance of Certification modules and the HIV Practice Improvement Module is encouraged as part of professional development for physicians who focus on HIV medicine.

beginning 36 months after certification or recertification to continue to be considered a qualified HIV physician.

Non-physician providers

Patient management experience and HIV-related continuing education also should be used to identify qualified nurse practitioners, physician assistants and nurse midwives who provide HIV primary care.

Other Considerations

<p>Areas of Lower HIV Prevalence *</p> <p>*Such as 6.8 cases per 100,000 or less</p> <p>Prevalence Rates available at: http://gis.cdc.gov/GRASP/NCHHSTPAtlas/main.html</p>	<p>The criteria above should NOT be used to exclude physicians from providing HIV care in areas with limited HIV workforce capacity. In communities or geographic areas where no physicians meet the criteria, medical providers are encouraged to develop a consultative relationship with a qualified adult or pediatric HIV/ID physician.</p>
<p>Pediatric/Adolescent Patient Management</p>	<p>The success of interventions to prevent perinatal HIV transmission has dramatically reduced the number of pediatric HIV cases in the U.S. However, managing pediatric and adolescent patients with HIV also requires appropriate expertise. In areas of low prevalence, less experienced physicians are encouraged to develop a consultative relationship as described above.</p>

Resources for Identifying HIV Medical Providers

- The Ryan White Program funds clinical sites to provide comprehensive HIV care in communities across the country. A directory of Ryan White-funded programs is available from the Health Resources and Services Administration website at <http://hab.hrsa.gov/>.
- The National HIV/AIDS Clinical Consultation Center supports a warmline for HIV providers available at 1-800-933-3413 or <http://www.nccc.ucsf.edu>.
- HIVMA maintains an online directory of members accepting new patients available at www.hivma.org. HIVMA does not credential or certify HIV medical providers.

HIV-related Standards of Care and Resources

- Federal HIV treatment guidelines on a range of topics, including antiretroviral treatment and the prevention and treatment of opportunistic infections are available at <http://aidsinfo.nih.gov/guidelines>.
- HIVMA guidelines on primary care and chronic kidney disease are available at www.hivma.org.
- HIV-related quality measures have been approved by the National Quality Forum (http://www.qualityforum.org/Measures_List.aspx) and included in Medicare’s Physician Quality Reporting System and Electronic Health Record Meaningful Use programs as well as the Initial Core Set of Health Quality Measures for Medicaid-Eligible Adults. Email info@hivma.org for additional information.

- A highly effective model for delivering HIV care has been developed by the Ryan White Program, the Department of Veterans Affairs and others. The model is detailed in *Essential Components of Effective HIV Care: A Policy Paper of the HIV Medicine Association of the Infectious Diseases Society of America and the Ryan White Medical Providers Coalition* (available online at www.hivma.org or by emailing info@hivma.org).

HIV Provider Experience and Patient Outcomes: Selected References

AIDS mortality rates lower at sites with HIV experience. *AIDS Alert* 1999;14(11):129-30.

Bach PB, Calhoun EA, Bennett CL. The relation between physician experience and patterns of care for patients with AIDS-related *Pneumocystis carinii* pneumonia: results from a survey of 1,500 physicians in the United States. *Chest* 1999;115(6):1563-9.

Brosgart C, et al. Community patterns of care for HIV disease: experience makes a difference. *Proceedings of Int Conf AIDS* 1998;12:1143-44.

Brosgart CL, et al. Clinical experience and choice of drug therapy for human immunodeficiency virus disease. *Clin Infect Dis* 1999;28(1):14-22.

Hecht FM, et al. Optimizing care for persons with HIV infection. Society of General Internal Medicine AIDS Task Force. *Ann Intern Med* 1999;131(2):136-43.

Horberg, et al. Influence of provider experience on antiretroviral adherence and viral suppression. *HIV AIDS (Auckl)* 2012;4:125-133.

Kitahata MM, Van Rompaey SE, Shields AW. Physician experience in the care of HIV-infected persons is associated with earlier adoption of new antiretroviral therapy. *J Acquir Immune Defic Syndr* 2000;24(2):106-14.

Kitahata MM, et al. Physicians experience with the acquired immunodeficiency syndrome as a factor in patients' survival. *N Engl J Med* 1996;334(11):701-706.

Laine C, et al. The relationship of clinic experience with advanced HIV and survival of women. *AIDS* 1998;12(4):417-424.

Landon, BE, et al. Physician specialization and the quality of care for human immunodeficiency virus infection. *Arch Intern Med* 2005;165:1133-1139.

Landon, BE, et al. Physician specialization and antiretroviral therapy for HIV: adoption and use in a national probability sample of persons infected with HIV. *J Gen Intern Med* 2003;18:233-241.

Markson LE, et al. Repeated emergency department use by HIV-infected persons: effect of clinic accessibility and expertise in HIV care. *J Acquir Immune Defic Syndr* 1998;17(1):35-41.

Moore RD, et al. Improvement in the health of HIV-infected persons in care: reducing disparities. *Clin Infect Dis* 2012;55(9):1242-51.

Shapiro M, et al. Variations in the care of HIV-infected adults in the United States. *JAMA* 1999;281:2305-2315.

Rackal, JM, et al. Provider training and experience for people living with HIV/AIDS. *Cochrane Database Syst Rev.* 2 2011;15(6):CD003938.

Stone VE, et al. Relation of physician specialty and HIV/AIDS experience to choice of guideline-recommended antiretroviral therapy. *J Gen Intern Med* 2001;16: 360-368.

Willard CL, Liljestrang P, Goldschmidt RH, Grumbach K. Is experience with human immunodeficiency virus disease related to clinical practice? A survey of rural primary care physicians. *Arch Fam Med* 1999; 8(6):502-8.

Wilson IB, Landon BE, Hirschhorn LR, et al. Quality of HIV Care Provided by Nurse Practitioners, Physician Assistants, and Physicians. *Ann Intern Med* 2005; 143(10): 729-36.